

Nagy, Örs; András, Szilárd

Approximated Poncelet configurations

formal überarbeitete Version der Originalveröffentlichung in:

formally revised edition of the original source in:

Teaching Mathematics and Computer Science 13 (2015) 2, S. 163-176



Bitte verwenden Sie in der Quellenangabe folgende URN oder DOI /

Please use the following URN or DOI for reference:

urn:nbn:de:0111-pedocs-121915

10.25656/01:12191

<https://nbn-resolving.org/urn:nbn:de:0111-pedocs-121915>

<https://doi.org/10.25656/01:12191>

Nutzungsbedingungen

Dieses Dokument steht unter folgender Creative Commons-Lizenz: <http://creativecommons.org/licenses/by-nc-sa/4.0/deed.de> - Sie dürfen das Werk bzw. den Inhalt unter folgenden Bedingungen vervielfältigen, verbreiten und öffentlich zugänglich machen sowie Abwandlungen und Bearbeitungen des Werkes bzw. Inhaltes anfertigen: Sie müssen den Namen des Autors/Rechteinhabers in der von ihm festgelegten Weise nennen. Dieses Werk bzw. der Inhalt darf nicht für kommerzielle Zwecke verwendet werden. Die neu entstandenen Werke bzw. Inhalte dürfen nur unter Verwendung von Lizenzbedingungen weitergegeben werden, die mit denen dieses Lizenzvertrages identisch oder vergleichbar sind.

Mit der Verwendung dieses Dokuments erkennen Sie die Nutzungsbedingungen an.

Terms of use

This document is published under following Creative Commons-License: <http://creativecommons.org/licenses/by-nc-sa/4.0/deed.en> - You may copy, distribute and transmit, adapt or exhibit the work in the public and alter, transform or change this work as long as you attribute the work in the manner specified by the author or licensor. You are not allowed to make commercial use of the work. If you alter, transform, or change this work in any way, you may distribute the resulting work only under this or a comparable license.

By using this particular document, you accept the above-stated conditions of use.



Kontakt / Contact:

peDOCS

DIPF | Leibniz-Institut für Bildungsforschung und Bildungsinformation

Informationszentrum (IZ) Bildung

E-Mail: pedocs@dipf.de

Internet: www.pedocs.de

Mitglied der


Leibniz-Gemeinschaft

Approximated Poncelet configurations

Örs Nagy¹, Szilárd András²
Babeş-Bolyai University, Cluj Napoca, Romania

Motto: *A picture is worth a thousand words*

ABSTRACT. In this short note we present the approximate construction of closed Poncelet configurations using the simulation of a mathematical pendulum. Although the idea goes back to the work of Jacobi ([16]), only the use of modern computer technologies assures the success of the construction. We present also some remarks on using such problems in project based university courses and we present a Matlab program able to produce animated Poncelet configurations with given period. In the same spirit we construct Steiner configurations and we give a few teaching oriented refinement on the Poncelet grid theorem.

KEYWORDS: mathematical pendulum, computer simulation, Poncelet porism, Steiner porism, Poncelet grid

ZDM SUBJECT CLASSIFICATION: U74, U75, G74, G75

MATHEMATICAL SUBJECT CLASSIFICATION: 68U20, 68U05, 97R60

Introduction

Consider two circles (or conics in the general case) Γ and γ . Starting from the point $A_0 \in \Gamma$ draw a tangent to γ which intersects Γ for the second time in A_1 . Repeating this construction we can define the sequence $(A_n)_{n \geq 0}$, where $A_k \in \Gamma$, $\forall k \geq 0$ and $A_k A_{k+1}$ is tangent to γ for all $k \geq 0$. This construction is called the Poncelet construction.

Let's recall Poncelet's famous closure theorem:

Theorem 1. ([18],[14],[10]) *If the sequence $(A_n)_{n \geq 0}$ from the Poncelet construction is periodic with period k for some point $A_0 \in \Gamma$, then it is periodic for all $A_0 \in \Gamma$.*

This theorem shows that the appearance of a closed Poncelet construction is determined by the two conics and their mutual position, so we can call (Γ, γ) a k -Poncelet configuration if the Poncelet construction has period k for all $A_0 \in \Gamma$. Results concerning the characterization of these configurations were established by Cayley in 1853 (see [5] or [10]) and recently by Dominique Hulin in 2007 (see [15]). Our intention was to give a computer algorithm (or a numerical method) for the construction of a Poncelet configurations with given period k in order to produce educational applets, animations. We have to mention that such configurations, animations are not available in dynamic geometric softwares or at web resources for $k \geq 5$. We have found a Java applet which generates animations for Poncelet's porism (see [21]) in the special case when Γ is a circle, γ is an ellipse and they have a common center of symmetry. For our purpose neither the Cayley conditions nor the Hulin decomposition proved to be useful in order to obtain an acceptable accuracy from the numerical approximations. Our approach relies on Jacobi's proof of the Poncelet theorem (see [16]) and uses the following property:

¹Email address: ors_nagy@yahoo.com

²Email address: andraszka@yahoo.com




Figure 1: Closed Poncelet construction with period $k = 5$

Theorem 2. Denote by $\Gamma = \partial\mathcal{C}(O, l)$ the circle of radius l obtained as the orbit of a mathematical pendulum with period T . Denote by A_j the position of the pendulum at the moment $j\frac{nT}{k}$ with $0 \leq j \leq k$. The lines $A_0A_1, A_1A_2, \dots, A_{k-1}A_k$ are tangents to a circle γ and the pair (Γ, γ) is a k -Poncelet configuration.

The above property shows that in order to produce Poncelet configurations we need to simulate the motion of a mathematical pendulum, to calculate the period T , to determine the coordinates of the points A_0, A_1, \dots, A_{k-1} and to calculate the coordinates of the center and the radius of the inner circle γ . Our Matlab program performs these steps and can be founded (together with some png animations) at

<http://www.math.ubbcluj.ro/~andrasz/poncelet/Animations.html>




Figure 2: Poncelet configurations with $k = 13, n = 6$ and $k = 15, n = 4$

If we replace the lines A_iA_{i+1} in the Poncelet theorem with circles \mathcal{C}_i for each $i \in \{0, 1, \dots, k-1\}$ such that \mathcal{C}_i and \mathcal{C}_{i+1} are tangent (in T_i) for $0 \leq i \leq k-1$ (where \mathcal{C}_k is \mathcal{C}_0) and all the circles \mathcal{C}_i are tangent to Γ and γ we obtain the Steiner theorem. The Steiner

theorem can be reduced to the Poncelet theorem if we observe that the loci of centers of the circles which are tangents to Γ and γ is an ellipse and the tangency points of T_i are on a fixed circle (see figure 3). An other proof of the Steiner theorem uses the fact that there exists an inversion which transforms Γ and γ into concentric circles. This idea can be used to generate Steiner configurations by constructing a corresponding Steiner configuration (with fixed n and k) for concentric circles and applying an inversion.




Figure 3: Steiner configurations with $k = 5$, $n = 1$ and $k = 5$, $n = 2$

If we take a closed look to the first case of figure 3 we can observe that the circles $\mathcal{C}_0, \mathcal{C}_1, \dots, \mathcal{C}_{k-1}$ may have intersection points (that are different from the tangency points). If Γ and γ are concentric circles than these intersection points are on some circles, hence this property remains true for the general case (see figure 4).




Figure 4: Additional properties of the Steiner configurations

A similar property of the Poncelet configurations is the recently discovered Poncelet grid (see [20]). On figure 5 we can observe that the set of all intersection points determined by two of the sides $A_0A_1, A_1A_2, \dots, A_{k-1}A_0$ can be partitioned such that each class of the

partition contains exactly k points and the points belonging to a class of the partition are moving on an ellipse when A_0 moves along Γ .




Figure 5: The ellipses containing the Poncelet grid

Moreover we can observe the following property

Theorem 3. *Consider a k -Poncelet configuration (Γ, γ) . If the points of a Poncelet grid are contained on the ellipses E_1, E_2, \dots, E_m and the points on E_j are labeled sequentially $X_{j,1}, X_{j,2}, \dots, X_{j,k}$ then for each $j \in \{1, 2, \dots, m\}$ there exists an ellipse³ which is tangent to the lines $X_{j1}X_{j,1+v}, X_{j2}X_{j,2+v}, \dots, X_{jk}X_{j,k+v}$.*

Remark 1. *This property is illustrated on figure 6 and shows that from a Poncelet configuration we can obtain infinitely many nested Poncelet grids belonging to Poncelet configurations with the same period k . The above theorem is mainly contained in [20] (theorem 1.1) but we think that it is useful to specify (especially for teaching reasons) that there are several Poncelet polygons with the same set of vertices. This is not clarified in [20] and the figures therein do not contain all the polygons.*

Remarks and teaching experience

We used the Poncelet closure theorem in our teaching activity at different levels. We worked with highschool students in several summer camps on the understanding of the elementary proofs. We have to mention that a completely elementary proof can be founded in the book of Sharigin ([19]). We included this theorem into an undergraduate geometry

³in the general case instead of ellipses we have conics




Figure 6: Additional properties of the Poncelet grid

course for university students in the first year, we used the Poncelet porism problem as an individual project subject for computer science students, we used the connection between the mathematical pendulum and the Poncelet theorem in a course on dynamical systems for computer science students and the study of new properties related to the Poncelet theorem (see [14], [3],[8], [20], [22]). The need of high quality visualization appeared at all these levels while it is almost impossible to draw or to construct exact figures if the period of construction k satisfies $k \geq 6$. As Howard Crosby wrote "A wisely chosen illustration is almost essential to fasten the truth upon the ordinary mind, and no teacher can afford to neglect this part of his preparation." In order to fulfill the necessity of a "wisely chosen illustration" we used our own figures. But at highschool level the proofs and our figures were not convincing enough (due to their complexity), the students understood the theorem but they were unable to construct their own Poncelet configuration and this lead to a serious frustration. Some very probable roots of such a frustration were formulated by S. Papert: "Better learning will not come from finding better ways for the teacher to instruct, but from giving the learner better opportunities to construct." and also by Kurt Levin: "If you want to truly understand something, try to change it." Unfortunately minor changes in the problem can lead to very hard problems, that they can't handle. We observed that at some of our students the initial frustration was transformed into a very deep motivation for further studies. We also have to point out that the use of animated Poncelet constructions helped a lot in the understanding of the Poncelet theorem and in the connection between the pendulum's motion and the Poncelet theorem. We had also a teaching activity where the students rediscovered the existence of the Poncelet grid and theorem 3 using the analogy between the Steiner and the Poncelet porism and constructing the corresponding animations. We can conclude that if a picture is worth a thousand words, then an animation (or simulation) is worth a thousand pictures.

On the other side these animations were not helpful in understanding the mathematical background and the proofs. This probably is connected with the ancient Chinese proverb "Tell me and I'll forget; show me and I may remember; involve me and I'll understand." We think that although the use of visualizations is indispensable we have to take care to avoid the situation when things are showed to students and they don't get involved.

Working with computer science students was a completely different experience because some of them get the problem as an individual project, so they had to develop a computer program which constructs Poncelet configurations. This framework assured that they got involved.

Proofs

For the sake of completeness we recall some well known facts about the mathematical pendulum and Jacobi's elliptic functions. For a few more details we recommend [1]. Consider a mathematical pendulum with length l and initial position characterized by $\varphi(0)$ and $\varphi'(0)$ (see figure 7). The motion of this pendulum is governed by the equation




Figure 7: The mathematical pendulum

$$\varphi'' + \frac{g}{l} \sin \varphi = 0.$$

Lemma 1. *The period of the pendulum can be expressed as*

$$T = 4\sqrt{\frac{l}{g}} E\left(\frac{\pi}{2}, \sin \frac{\varphi_0}{2}\right),$$

where

$$E(\varphi, k) = \int_0^\varphi \frac{dt}{\sqrt{1 - k^2 \sin^2 t}}$$

is the elliptic integral of the first kind.

Definition 1. *If $k \in (0, 1)$ and*

$$E(\varphi) = \int_0^\varphi \frac{dt}{\sqrt{1 - k^2 \sin^2 t}}$$

is the elliptic integral of the first kind, then with the inverse $z \rightarrow \text{Am}(z)$ of the function $\varphi \rightarrow E(\varphi)$ ($\text{Am}(z) = \varphi \Leftrightarrow z = E(\varphi)$) we can define Jacobi's elliptic functions:

$$\text{sn}(z) = \sin(\text{Am}(z));$$

$$\begin{aligned}\operatorname{cn}(z) &= \cos(\operatorname{Am}(z)); \\ \operatorname{dn}(z) &= \sqrt{1 - k^2 \sin^2 \operatorname{Am}(z)}.\end{aligned}$$

Lemma 2. *The solution of the Cauchy problem $\varphi''(t) = -k^2 \sin(\varphi(t))$ $\varphi(t_0) = \varphi_0$ and $\varphi'(t_0) = \varphi'_0$ is the function*

$$\varphi(t) = 2 \operatorname{Am}(\nu(t - t_0) + K_1), \quad (1)$$

where

$$\begin{aligned}K_1 &= \int_0^{\varphi_0/2} \frac{du}{\sqrt{1 - \rho^2 \sin^2(u)}}, \quad \rho = \frac{k}{\nu} \\ \nu &= \sqrt{\frac{1}{4}(\varphi'_0)^2 + k^2 \sin^2\left(\frac{\varphi_0}{2}\right)}.\end{aligned}$$

Proof of theorem 2 and 3. Consider $\Gamma = \partial\mathcal{C}(O, l)$ the orbit of the pendulum and $\mathcal{C}(I, r)$ the circle with center I and radius r . Let O be the origin and OI the Ox axis, so $\overrightarrow{OI} = \lambda \vec{i}$ where $|\lambda| + r < l$. If $Q_k \in \Gamma$, $k \in \{1, 2\}$ are two distinct points, then there exist $(\alpha_1, \alpha_2) \in \mathbb{R}^2$ with $\alpha_1 - \alpha_2 \notin 2\pi\mathbb{Z}$ and $\overrightarrow{OQ_k} = l(\vec{i} \cos \alpha_k + \vec{j} \sin \alpha_k)$. The equation of the line Q_1Q_2 is

$$L(x, y) := x \cos\left(\frac{\alpha_1 + \alpha_2}{2}\right) + y \sin\left(\frac{\alpha_1 + \alpha_2}{2}\right) - l \cos\left(\frac{\alpha_1 - \alpha_2}{2}\right) = 0. \quad (2)$$

The distance from $I(\lambda, 0)$ to the line Q_1Q_2 is $|L(\lambda, 0)|$, so Q_1Q_2 is tangent to the interior circle if and only if $L(\lambda, 0) = \varepsilon r$, where $\varepsilon \in \{-1, 1\}$. Hence the tangents to the interior circle can be characterized by the equation

$$\lambda \cos\left(\frac{\alpha_1 + \alpha_2}{2}\right) - l \cos\left(\frac{\alpha_1 - \alpha_2}{2}\right) - \varepsilon r = 0. \quad (3)$$

Denote by T the period of the pendulum, by k the period of the desired construction and by n the number of pendulum periods used for the construction. In addition for each $0 \leq j \leq k$ denote by A_j the position of the pendulum at the moment $t_j = j\tau$ with $\tau = \frac{n\pi}{k}$. In order to prove theorem 2 and 3 it is sufficient to prove that the lines A_jA_{j+v} , $0 \leq j \leq k$ are tangents of a fixed circle $\mathcal{C}(I, r)$. Moreover we prove that if $\Theta_p(t)$ is the solution of the pendulum's equation (with initial conditions $\varphi(t_0)$ and $\varphi'(t_0)$) at the moment $t - p\tau$ and $A_p(t)$ the position of the pendulum, then for all t the lines $A_p(t)A_{p+v}(t)$ are tangents⁴ to a fixed circle $\mathcal{C}_v(I, r)$. Due to lemma 2 for $\varphi_0 = t_0 = 0$ we have $\Theta_p(t) = \varphi(t - p\tau) = 2 \operatorname{Am}(\nu(t - p\tau))$, where $\nu = \frac{1}{2}|\dot{\varphi}_0|$.

According to (2) the equation of the line $A_p(t)A_{p+v}(t)$ is:

$$\begin{aligned}x \cos\left(\frac{\Theta_p(t) + \Theta_{p+v}(t)}{2}\right) + y \sin\left(\frac{\Theta_p(t) + \Theta_{p+v}(t)}{2}\right) - \\ - l \cos\left(\frac{\Theta_p(t) - \Theta_{p+v}(t)}{2}\right) = 0\end{aligned}$$

⁴This is a key element in designing the animations.

With the notations $s := \nu(t - p\tau)$ and $\delta = v\tau$ we have $\nu(t - (p+v)\tau) = s - \nu v\tau = s - \delta$, so

$$\begin{aligned} \cos\left(\frac{\Theta_p(t) + \Theta_{p+v}(t)}{2}\right) &= \cos(\text{Am}(s) + \text{Am}(s - \delta)) = \\ &= \text{cn}(s) \text{cn}(s - \delta) - \text{sn}(s) \text{sn}(s - \delta) \end{aligned} \quad (4)$$

and

$$\cos\left(\frac{\Theta_p(t) - \Theta_{p+v}(t)}{2}\right) = \text{cn}(s) \text{cn}(s - \delta) + \text{sn}(s) \text{sn}(s - \delta). \quad (5)$$

Due to (4) and (5) for the expression

$$\mathcal{S} := l \frac{\text{dn}(\delta) - 1}{\text{dn}(\delta) + 1} \cos\left(\frac{\Theta_p(t) + \Theta_{p+v}(t)}{2}\right) - l \cos\left(\frac{\Theta_p(t) - \Theta_{p+v}(t)}{2}\right)$$

we obtain

$$\mathcal{S} = -\frac{2l}{1 + \text{dn}(\delta)} (\text{cn}(s) \text{cn}(s - \delta) + \text{sn}(s) \text{sn}(s - \delta) \text{dn}(\delta)), \quad (6)$$

which leads to

$$\mathcal{S} = -\frac{2l \text{cn}(\delta)}{1 + \text{dn}(\delta)}. \quad (7)$$

This implies that the lines $A_p(t)A_{p+v}(t)$ are tangents to the circle $\mathcal{C}(I, r)$ for all $t \in \mathbb{R}$ if

$$\overrightarrow{OI} = l \frac{\text{dn}(\delta) - 1}{\text{dn}(\delta) + 1} \vec{i},$$

and

$$r = \frac{2l |\text{cn}(\delta)|}{1 + \text{dn}(\delta)}.$$

□

Remark 2. *This proof is in fact Jacobi's original proof (with somewhat modified notations) for $v = 1$. For $1 \leq v \leq [(k-1)/2]$ we obtain different polygons with the same vertices. If we denote by $X_1 X_2 \dots X_k$ the convex Poncelet polygon inscribed in Γ , then the points of the associated Poncelet grid can be obtained by constructing all diagonals of the polygon $X_1 X_2 \dots X_k$. Moreover if E_v is a Poncelet "gridline", then the intersection points $X_{j1} X_{j2}, \dots, X_{jk}$ generate an other Poncelet grid.*

Remark 3. *Using the same ideas as in the above proof we can show that the intersection points $A_i A_j \cap A_{i+v} A_{j+v}$ belong to an ellipse for fixed i, j and $1 \leq v \leq k$, so this approach represents an alternative proof for theorem 1.1. from [20].*

Concluding remarks

- The use of visualizations and animations is strongly recommended in the teaching of mathematics. Moreover it is helpful if the students can generate their own animations using some mathematical software (Matlab, Mathematica). In order to overcome this need the teacher training curricula must contain special courses on

the use of modern technology. The Poncelet and the Steiner theorem represents a very good teaching example in this direction because the configurations can not be constructed without a computer for arbitrary k .

- The parallel use of modern technologies (computers) and traditional methods/accessories is an imperative necessity of high quality inquiry based mathematical education. This implies that the classical classroom settings, the organization of activities must be completely restructured in order to fulfill this necessity and to improve performance.
- It would be interesting to generate also Zig-Zag and Ponzag configurations using the equivalences from [13] and [14].
- It would be helpful in many teaching situations to include the Poncelet and the Steiner porism into dynamic geometric softwares like Geonext, Geogebra, Cabri.

Acknowledgements

A part of our teaching activity was organized in the framework of the project DQME II⁵. The first author's research was partially supported by the Hungarian University Federation of Cluj Napoca.

References

- [1] András, Szilárd: *Dinamikus rendszerek*, Editura Didactică și Pedagogică, București, 2008
- [2] Bråting, Kajsa; Pejlar, Johanna: *Visualizations in mathematics*, Erkenntnis 68 (2008), no. 3, 345–358.
- [3] Burskii, V.P. and Zhedanov, A.S.: *Boundary-value problems for the string equation, the poncelet problem, and the Pell-Abel equation: links and relations*, Journal of Mathematical Sciences, 149(2008), 5, 1483-1487
- [4] A. Cayley⁶: *Note on the geometrical representation of the integral $\int dx \div \sqrt{(x+a)(x+b)(x+c)}$* , Philosophical Magazine, vol. V (1853), 99-102.
- [5] A. Cayley: *Note on the Porism of the in-and-circumscribed polygon*, Philosophical Magazine, vol. VI (1853), 376-377.
- [6] A. Cayley: *Correction of two theorems relating to the porism of the in-and-circumscribed polygon* Philosophical Magazine, vol. VI (1853), 376-377.
- [7] A. Cayley: *Developments on the porism of the in-and-circumscribed polygon*, Philosophical Magazine, vol. VII (1854), 339-345.

⁵Developing Quality in Mathematics Education, for more details see <http://www.dqime.uni-dortmund.de/>

⁶ For items [4]-[7] see papers 113, 115, 116, 128 in Collected Works of Cayley

- [8] Daepf U., Gorkin P. and Voss, K.: *Poncelet's theorem, Sendov's conjecture, and Blaschke products*, J. Math. Anal. Appl. (2009), doi:10.1016/j.jmaa.2009.09.058
- [9] Darboux, Gaston: *Principes de Géométrie Analytique*, Gauthier-Villars et Cie Éditeurs, Paris, 1917.
- [10] Flatto, Leopold: *Poncelet's Theorem*, American Mathematical Society, 2008
- [11] Giaquinto, M.: *Visual thinking in mathematics. An epistemological study*, Oxford University Press, Oxford, 2007.
- [12] Herr, Ulrike: *Über das Theorem von Poncelet*, Staatsexamensarbeit, Johannes Gutenberg Universität, Mainz, 2000.
- [13] Hráskó, András: *Geometria tételek a harmadrendű görbe csoporttulajdonságával összefüggésben*, PhD értekezés, Eötvös Loránd Tudományegyetem, Budapest, 2005
- [14] Hráskó, András: *Poncelet-type Problems, an Elementary Approach*, Elem. Math. 55(2000), 45-62
- [15] Hulin, Dominique: *Factorisation des polynômes de Poncelet*, Geom Dedicata (2007) 130:109-136
- [16] C.G.J. Jacobi: *Über die Anwendung der Elliptischen Transcendenten auf ein bekanntes Problem der Elementargeometrie*, Crelle Journal für die reine und angewandte Mathematik, 3 (1828), 376-389, and in: *Gesammelte Werke*, 278-293.
- [17] Lebesgue, Henri: *Les Coniques*, Gauthier-Villars Imprimeur-Éditeur, Paris, 1942.
- [18] Poncelet, Jean-Victor: *Traité des propriétés projectives des figures*, Paris, 1822.
- [19] Sharigin, I.M.: *Problems in plane geometry*, MIR Publishers, Moscow, 1989
- [20] Schwartz, Richard Evan: *The Poncelet grid*, Adv. Geom. 7 (2007), no. 2, 157-175
- [21] Schwartz, Richard Evan: Java Applet for Poncelet's porism, <http://www.math.brown.edu/~res/Java/App37/test1.html>
- [22] Tabachnikov, Serge; Levi, Mark: *The Poncelet grid and billiards in ellipses*, Amer. Math. Monthly 114 (2007), no. 10, 895-908.